

RESEARCH

Open Access

Approximating fixed points of amenable semigroup and infinite family of nonexpansive mappings and solving systems of variational inequalities and systems of equilibrium problems

Hossein Piri

Correspondence:
hossein_piri1979@yahoo.com
Department of Mathematics,
University of Bonab 55517-61167
Bonab, Iran

Abstract

We introduce an iterative scheme for finding a common element of the set of solutions for systems of equilibrium problems and systems of variational inequalities and the set of common fixed points for an infinite family and left amenable semigroup of nonexpansive mappings in Hilbert spaces. The results presented in this paper mainly extend and improved some well-known results in the literature.

Mathematics Subject Classification (2000): 47H09; 47H10; 47H20; 43A07; 47J25.

Keywords: common fixed point, strong convergence, amenable semigroup, explicit iterative, system of equilibrium problem.

1. Introduction

Let H be a real Hilbert space and let C be a nonempty closed convex subset of H .

Let $A: C \rightarrow H$ be a nonlinear mapping. The classical variational inequality problem is to find $x \in C$ such that

$$\langle Ax, y - x \rangle \geq 0, \quad \forall y \in C. \quad (1)$$

The set of solution of (1) is denoted by $VI(C, A)$, i.e.,

$$VI(C, A) = \{x \in C : \langle Ax, y - x \rangle \geq 0, \quad \forall y \in C\}. \quad (2)$$

Recall that the following definitions:

(1) A is called monotone if

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad \forall x, y \in C.$$

(2) A is called α -strongly monotone if there exists a positive constant α such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|x - y\|^2, \quad \forall x, y \in C.$$

(3) A is called μ -Lipschitzian if there exist a positive constant μ such that

$$\|Ax - Ay\| \leq \mu \|x - y\|, \quad \forall x, y \in C.$$

(4) A is called α -inverse strongly monotone, if there exists a positive real number $\alpha > 0$

such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad \forall x, y \in C.$$

It is obvious that any α -inverse strongly monotone mapping B is $\frac{1}{\alpha}$ -Lipschitzian.

(5) A mapping $T : C \rightarrow C$ is called nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. Next, we denote by $\text{Fix}(T)$ the set of fixed point of T .

(6) A mapping $f : C \rightarrow C$ is said to be contraction if there exists a coefficient $\alpha \in (0, 1)$ such that

$$\|f(x) - f(y)\| \leq \alpha \|x - y\|, \quad \forall x, y \in C.$$

(7) A set-valued mapping $U : H \rightarrow 2^H$ is called monotone if for all $x, y \in H, f \in Ux$ and $g \in Uy$ imply $\langle x - y, f - g \rangle \geq 0$.

(8) A monotone mapping $U : H \rightarrow 2^H$ is maximal if the graph $G(U)$ of U is not properly contained in the graph of any other monotone mapping.

It is known that a monotone mapping U is maximal if and only if for $(x, f) \in H \times H$, $\langle x - y, f - g \rangle \leq 0$ for every $(y, g) \in G(U)$ implies that $f \in Ux$. Let B be a monotone mapping of C into H and let $N_C x$ be the normal cone to C at $x \in C$, that is, $N_C x = \{y \in H : \langle x - z, y \rangle \leq 0, \forall z \in C\}$ and define

$$Ux = \begin{cases} Bx + N_C x, & x \in C, \\ \emptyset & x \notin C. \end{cases}$$

Then U is the maximal monotone and $0 \in Ux$ if and only if $x \in \text{VI}(C, B)$; see [1].

Let F be a bi-function of $C \times C$ into \mathbb{R} , where \mathbb{R} is the set of real numbers. The equilibrium problem for $F : C \times C \rightarrow \mathbb{R}$ is to determine its equilibrium points, i.e the set

$$\text{EP}(F) = \{x \in C : F(x, y) \geq 0, \forall y \in C\}.$$

Let $\mathcal{J} = \{F_i\}_{i \in I}$ be a family of bi-functions from $C \times C$ into \mathbb{R} . The system of equilibrium problems for $\mathcal{J} = \{F_i\}_{i \in I}$ is to determine common equilibrium points for $\mathcal{J} = \{F_i\}_{i \in I}$, i.e the set

$$\text{EP}(\mathcal{J}) = \{x \in C : F_i(x, y) \geq 0, \forall y \in C, \forall i \in I\}. \quad (3)$$

Numerous problems in physics, optimization, and economics reduce into finding some element of $\text{EP}(F)$. Some method have been proposed to solve the equilibrium problem; see, for instance [2-5]. The formulation (3), extend this formalism to systems of such problems, covering in particular various forms of feasibility problems [6,7].

Given any $r > 0$ the operator $J_r^F : H \rightarrow C$ defined by

$$J_r^F(x) = \{z \in C : F(z, \gamma) + \frac{1}{r} \langle \gamma - z, z - x \rangle \geq 0, \forall \gamma \in C\},$$

is called the resolvent of F , see [3]. It is shown [3] that under suitable hypotheses on F (to be stated precisely in Sect. 2), $J_r^F : H \rightarrow C$ is single-valued and firmly nonexpansive and satisfies

$$\text{Fix}(J_r^F) = \text{EP}(F), \quad \forall r > 0.$$

Using this result, in 2007, Yao et al. [8], proposed the following explicit scheme with respect to W -mappings for an infinite family of nonexpansive mappings:

$$x_{n+1} = \alpha_n f(x_n) + \beta_n x_n + \gamma_n W_n J_{r_n}^F x_n \quad (4)$$

They proved that if the sequences $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ and $\{r_n\}$ of parameters satisfy appropriate conditions, then, the sequences $\{x_n\}$ and $\{J_{r_n}^F x_n\}$ both converge strongly to the unique $x^* \in \bigcap_{i=1}^{\infty} \text{Fix}(T_i) \cap \text{EP}(F)$, where $x^* \in P_{\bigcap_{i=1}^{\infty} \text{Fix}(T_i) \cap \text{EP}(F)} f(x^*)$. Their results extend and improve the corresponding results announced by Combettes and Hirstoaga [3] and Takahashi and Takahashi [5].

Very recently, Jitpeera et al. [9], introduced the iterative scheme based on viscosity and Cesàro mean

$$\begin{cases} \phi(u_n, \gamma) + \phi(\gamma) - \phi(u_n) + \frac{1}{r_n} \langle \gamma - u_n, u_n - x_n \rangle \geq 0, & \forall \gamma \in C, \\ \gamma_n = \delta_n u_n + (1 - \delta_n) P_C(u_n - \lambda_n B u_n), \\ x_{n+1} = \alpha_n \gamma f(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n A) \frac{1}{n+1} \sum_{i=0}^n T^i \gamma_n, & \forall n \geq 0, \end{cases}$$

where $B : C \rightarrow H$ is β -inverse strongly monotone, $\phi : C \rightarrow \mathbb{R} \cup \{\infty\}$ is a proper lower semi-continuous and convex function, $T^i : C \rightarrow C$ is a nonexpansive mapping for all $i = 1, 2, \dots, n$, $\{\alpha_n\}$, $\{\beta_n\}$, $\{\delta_n\} \subset (0, 1)$, $\{\lambda_n\} \subset (0, 2\beta)$ and $\{r_n\} \subset (0, \infty)$ satisfy the following conditions

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (ii) $\lim_{n \rightarrow \infty} \delta_n = 0$
- (iii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.
- (iv) $\{\lambda_n\} \subset [a, b] \subset (0, 2\beta)$ and $\liminf_{n \rightarrow \infty} |\lambda_{n+1} - \lambda_n| = 0$,
- (v) $\liminf_{n \rightarrow \infty} r_n > 0$ and $\liminf_{n \rightarrow \infty} |r_{n+1} - r_n| = 0$.

They show that if $\theta = \bigcap_{i=1}^n \text{Fix}(T^i) \cap \text{VI}(C, B) \cap \text{MEP}(\phi, \varphi)$ is nonempty, then the sequence $\{x_n\}$ converges strongly to the $z = P_{\theta}(I - A + \gamma f)z$ which is the unique solution of the variational inequality

$$\langle (\gamma f - A)z, x - z \rangle \leq 0 \quad \forall x \in \theta.$$

In this paper, motivated and inspired by Yao et al. [8,10-15], Lau et al. [16], Jitpeera et al. [9], Kangtunyakarn [17] and Kim [18], Atsushiba and Takahashi [19], Saeidi [20], Piri [21-23] and Piri and Badali [24], we introduce the following iterative scheme for finding a common element of the set of solutions for a system of equilibrium problems

$\mathcal{J} = \{F_k : k = 1, 2, 3, \dots, M\}$ for a family $\mathcal{J} = \{F_k : k = 1, 2, 3, \dots, M\}$ of equilibrium bifunctions, systems of variational inequalities, the set of common fixed points for an infinite family $\psi = \{T_i, i = 1, 2, \dots\}$ of nonexpansive mappings and a left amenable semigroup $\phi = \{T_t : t \in S\}$ of nonexpansive mappings, with respect to W -mappings and a left regular sequence $\{\mu_n\}$ of means defined on an appropriate space of bounded real-valued functions of the semigroup

$$\begin{cases} z_n = J_{r_M, n}^{F_M} \dots J_{r_2, n}^{F_2} J_{r_1, n}^{F_1} x_n, \\ \gamma_n = \eta_n P_C(z_n - \zeta_n A z_n) + (1 - \eta_n) P_C(z_n - \delta_n B z_n), \\ x_{n+1} = \alpha_n f(T_{\mu_n} W_n \gamma_n) + \beta_n x_n + \gamma_n T_{\mu_n} W_n \gamma_n, \quad n \geq 1, \end{cases} \quad (5)$$

where $A: C \rightarrow H$ be β -inverse monotone map and $B: C \rightarrow H$ be δ -inverse monotone map. We prove that under mild assumptions on parameters like that in Yao et al. [8], the sequences $\{x_n\}$ and $\{J_{r_{k_n}, n}^{F_k} x_n\}_{k=1}^M$ converge strongly to $x^* \in \mathcal{F} = \bigcap_{i=1}^{\infty} \text{Fix}(T_i) \cap \text{Fix}(\varphi) \cap \text{EP}(\mathcal{J}) \cap \text{VI}(C, A) \cap \text{VI}(C, B)$, where $x^* = P_{\mathcal{F}}(x^*)$.

Compared to the similar works, our results have the merit of studying the solutions of systems of equilibrium problems, systems of variational inequalities and fixed point problems of amenable semigroup of nonexpansive mappings. Consequence for nonnegative integer numbers is also presented.

2. Preliminaries

Let S be a semigroup and let $B(S)$ be the space of all bounded real valued functions defined on S with supremum norm. For $s \in S$ and $f \in B(S)$, we define elements $l_s f$ and $r_s f$ in $B(S)$ by

$$(l_s f)(t) = f(st), \quad (r_s f)(t) = f(ts), \quad \forall t \in S.$$

Let X be a subspace of $B(S)$ containing 1 and let X^* be its topological dual. An element μ of X^* is said to be a mean on X if $\|\mu\| = \mu(1) = 1$. We often write $\mu_t(f(t))$ instead of $\mu(f)$ for $\mu \in X^*$ and $f \in X$. Let X be left invariant (respectively right invariant), i.e., $l_s(X) \subset X$ (respectively $r_s(X) \subset X$) for each $s \in S$. A mean μ on X is said to be left invariant (respectively right invariant) if $\mu(l_s f) = \mu(f)$ (respectively $\mu(r_s f) = \mu(f)$) for each $s \in S$ and $f \in X$. X is said to be left (respectively right) amenable if X has a left (respectively right) invariant mean. X is amenable if X is both left and right amenable. As is well known, $B(S)$ is amenable when S is a commutative semigroup, see [25]. A net $\{\mu_\alpha\}$ of means on X is said to be strongly left regular if

$$\lim_{\alpha} \|l_s^* \mu_\alpha - \mu_\alpha\| = 0,$$

for each $s \in S$, where l_s^* is the adjoint operator of l_s .

Let S be a semigroup and let C be a nonempty closed and convex subset of a reflexive Banach space E . A family $\phi = \{T_t : t \in S\}$ of mapping from C into itself is said to be a nonexpansive semigroup on C if T_t is nonexpansive and $T_{ts} = T_t T_s$ for each $t, s \in S$. By $\text{Fix}(\phi)$ we denote the set of common fixed points of ϕ , i.e.

$$\text{Fix}(\varphi) = \bigcap_{t \in S} \{x \in C : T_t x = x\}.$$

Lemma 2.1. [25] Let S be a semigroup and C be a nonempty closed convex subset of a reflexive Banach space E . Let $\phi = \{T_t : t \in S\}$ be a nonexpansive semigroup on H such that $\{T_t x : t \in S\}$ is bounded for some $x \in C$, let X be a subspace of $B(S)$ such that $1 \in X$ and the mapping $t \rightarrow \langle T_t x, y^* \rangle$ is an element of X for each $x \in C$ and $y^* \in E^*$, and μ is a mean on X . If we write $T_\mu x$ instead of $\int T_t x d\mu(t)$, then the followings hold.

- (i) T_μ is nonexpansive mapping from C into C .
- (ii) $T_\mu x = x$ for each $x \in \text{Fix}(\phi)$.
- (iii) $T_\mu x \in \overline{\text{co}}\{T_t x : t \in S\}$ for each $x \in C$.

Let C be a nonempty subset of a Hilbert space H and $T : C \rightarrow H$ a mapping. Then T is said to be demiclosed at $v \in H$ if, for any sequence $\{x_n\}$ in C , the following implication holds:

$$x_n \rightarrow u \in C, \quad Tx_n \rightarrow v \quad \text{imply} \quad Tu = v,$$

where \rightarrow (respectively \rightharpoonup) denotes strong (respectively weak) convergence.

Lemma 2.2. [26] Let C be a nonempty closed convex subset of a Hilbert space H and suppose that $T : C \rightarrow H$ is nonexpansive. then, the mapping $I - T$ is demiclosed at zero.

Lemma 2.3. [27] For a given $x \in H$, $y \in C$,

$$y = P_C x \Leftrightarrow \langle y - x, z - y \rangle \geq 0, \quad \forall z \in C.$$

It is well known that P_C is a firmly nonexpansive mapping of H onto C and satisfies

$$\|P_C x - P_C y\|^2 \leq \langle P_C x - P_C y, x - y \rangle, \quad \forall x, y \in H. \quad (6)$$

Moreover, P_C is characterized by the following properties: $P_C x \in C$ and for all $x \in H$, $y \in C$,

$$\langle x - P_C x, y - P_C x \rangle \leq 0. \quad (7)$$

It is easy to see that (7) is equivalent to the following inequality

$$\|x - y\|^2 \geq \|x - P_C x\|^2 + \|y - P_C x\|^2. \quad (8)$$

Using Lemma 2.3, one can see that the variational inequality (1) is equivalent to a fixed point problem. It is easy to see that the following is true:

$$u \in \text{VI}(C, A) \Leftrightarrow u = P_C(u - \lambda Au), \quad \lambda > 0. \quad (9)$$

Lemma 2.4. [28] Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space E and let $\{\alpha_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$. Suppose $x_{n+1} = \alpha_n x_n + (1 - \alpha_n)y_n$ for all integers $n \geq 0$ and

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 1.$$

Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.

Let $F : C \times C \rightarrow \mathbb{R}$ be a bi-function. Given any $r > 0$, the operator $J_r^F : H \rightarrow C$ defined by

$$J_r^F x = \left\{ z \in C : F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\}$$

is called the resolvent of F , see [3]. The equilibrium problem for F is to determine its equilibrium points, i.e., the set

$$\text{EP}(F) = \{x \in C : F(x, y) \geq 0, \forall y \in C\}.$$

Let $\mathcal{J} = \{F_i\}_{i \in I}$ be a family of bi-functions from $C \times C$ into \mathbb{R} . The system of equilibrium problems for \mathcal{J} is to determine common equilibrium points for $\mathcal{J} = \{F_i\}_{i \in I}$, i.e., the set

$$\text{EP}(\mathcal{J}) = \{x \in C : F_i(x, y) \geq 0, \forall y \in C, \forall i \in I\}.$$

Lemma 2.5. [3] *Let C be a nonempty closed convex subset of H and $F : C \times C \rightarrow \mathbb{R}$ satisfy*

- (A₁) $F(x, x) = 0$ for all $x \in C$,
- (A₂) F is monotone, i.e, $F(x, y) + F(y, x) \leq 0$ for all $x, y \in C$,
- (A₃) for all $x, y, z \in C$, $\lim_{t \rightarrow 0} F(tz + (1-t)x, y) \leq F(x, y)$,
- (A₄) for all $x \in C$, $y \rightarrow F(x, y)$ is convex and lower semi-continuous.

Given $r > 0$, define the operator $J_r^F : H \rightarrow C$, the resolvent of F , by

$$J_r^F(x) = \{z \in C : F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C\}.$$

Then,

- (1) J_r^F is single valued,
- (2) J_r^F is firmly nonexpansive, i.e, $\|J_r^F x - J_r^F y\|^2 \leq \langle J_r^F x - J_r^F y, x - y \rangle$ for all $x, y \in H$,
- (3) $\text{Fix}(J_r^F) = \text{EP}(F)$,
- (4) $\text{EP}(F)$ is closed and convex.

Let T_1, T_2, \dots be an infinite family of mappings of C into itself and let $\lambda_1, \lambda_2, \dots$ be a real numbers such that $0 \leq \lambda_i < 1$ for every $i \in \mathbb{N}$. For any $n \in \mathbb{N}$, define a mapping W_n of C into C as follows:

$$\begin{aligned} U_{n,n+1} &= I, \\ U_{n,n} &= \lambda_n T_n U_{n,n+1} + (1 - \lambda_n) I, \\ U_{n,n-1} &= \lambda_{n-1} T_{n-1} U_{n,n} + (1 - \lambda_{n-1}) I, \\ &\vdots \\ U_{n,k} &= \lambda_k T_k U_{n,k+1} + (1 - \lambda_k) I, \\ U_{n,k-1} &= \lambda_{k-1} T_{k-1} U_{n,k} + (1 - \lambda_{k-1}) I, \\ &\vdots \\ U_{n,2} &= \lambda_2 T_2 U_{n,3} + (1 - \lambda_2) I, \\ W_n = U_{n,1} &= \lambda_1 T_1 U_{n,2} + (1 - \lambda_1) I. \end{aligned} \tag{10}$$

Such a mapping W_n is called the W -mapping generated by T_1, T_2, \dots, T_n and $\lambda_1, \lambda_2, \dots, \lambda_n$.

Lemma 2.6. [29] *Let C be a nonempty closed convex subset of a Hilbert space H , $\{T_i : C \rightarrow C\}$ be an infinite family of nonexpansive mappings with $\bigcap_{i=1}^{\infty} \text{Fix}(T_i) \neq \emptyset$, $\{\lambda_i\}$ be a real sequence such that $0 < \lambda_i \leq b < 1, \forall i \geq 1$. Then*

- (1) W_n is nonexpansive and $\text{Fix}(W_n) = \bigcap_{i=1}^n \text{Fix}(T_i)$ for each $n \geq 1$,

(2) for each $x \in C$ and for each positive integer j , the limit $\lim_{n \rightarrow \infty} U_{n,j}$ exists.

(3) The mapping $W : C \rightarrow C$ defined by

$$Wx := \lim_{n \rightarrow \infty} W_n x = \lim_{n \rightarrow \infty} U_{n,1} x, \quad \forall x \in C,$$

is a nonexpansive mapping satisfying $\text{Fix}(W) = \bigcap_{i=1}^{\infty} \text{Fix}(T_i)$ and it is called the W -mapping generated by T_1, T_2, \dots and $\lambda_1, \lambda_2, \dots$.

Lemma 2.7. [30] Let C be a nonempty closed convex subset of a Hilbert space H , $\{T_i : C \rightarrow C\}$ be a countable family of nonexpansive mappings with $\bigcap_{i=1}^{\infty} \text{Fix}(T_i) \neq \emptyset$, $\{\lambda_i\}$ be a real sequence such that $0 < \lambda_i \leq b < 1$, $\forall i \geq 1$. If D is any bounded subset of C , then

$$\lim_{n \rightarrow \infty} \sup_{x \in D} \|Wx - W_n x\| = 0.$$

Lemma 2.8. [31] Let $\{a_n\}$ be a sequence of nonnegative real numbers such that

$$a_{n+1} \leq (1 - b_n)a_n + b_n c_n, \quad n \geq 0,$$

where $\{b_n\}$ and $\{c_n\}$ are sequences of real numbers satisfying the following conditions:

- (i) $\{b_n\} \subset [0, 1]$, $\sum_{n=0}^{\infty} b_n = \infty$,
- (ii) either $\limsup_{n \rightarrow \infty} c_n \leq 0$ or $\sum_{n=0}^{\infty} |b_n c_n| < \infty$.

Then, $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2.9. [32] Let $(E, \langle \cdot, \cdot \rangle)$ be an inner product space. Then for all $x, y, z \in E$ and $\alpha, \beta, \gamma \in [0, 1]$ such that $\alpha + \beta + \gamma = 1$, we have

$$\begin{aligned} \|\alpha x + \beta y + \gamma z\|^2 &= \alpha \|x\|^2 + \beta \|y\|^2 + \gamma \|z\|^2 \\ &\quad - \alpha \beta \|x - y\|^2 - \alpha \gamma \|x - z\|^2 - \beta \gamma \|y - z\|^2. \end{aligned}$$

Notation Throughout the rest of this paper the open ball of radius r centered at 0 is denoted by B_r . For a subset A of H we denote by $\overline{\text{co}}A$ the closed convex hull of A . For $\epsilon > 0$ and a mapping $T : D \rightarrow H$, we let $F_{\epsilon}(T; D)$ be the set of ϵ -approximate fixed points of T , i.e., $F_{\epsilon}(T; D) = \{x \in D : \|x - Tx\| \leq \epsilon\}$. Weak convergence is denoted by \rightharpoonup and strong convergence is denoted by \rightarrow .

3. Strong convergence

Theorem 3.1. Let C be a nonempty closed convex subset of a real Hilbert space H , $A : C \rightarrow H$ a β -inverse strongly monotone, $B : C \rightarrow H$ a γ -inverse strongly monotone, S a semigroup and $\phi = \{T_t : t \in S\}$ be a nonexpansive semigroup from C into C such that $\text{Fix}(\phi) = \bigcap_{t \in S} \text{Fix}(T_t) \neq \emptyset$. Let X be a left invariant subspace of $B(S)$ such that $1 \in X$, and the function $t \rightarrow \langle T_t x, y \rangle$ is an element of X for each $x \in C$ and $y \in H$, $\{\mu_n\}$ a left regular sequence of means on X such that $\lim_{n \rightarrow \infty} \|\mu_{n+1} - \mu_n\| = 0$. Let $\mathcal{J} = \{F_k : k = 1, 2, \dots, M\}$ be a finite family of bi-functions from $C \times C$ into \mathbb{R} which satisfy (A_1) -(A_4) and $\{T_i\}_{i=1}^{\infty}$ an infinite family of nonexpansive mappings of C into C such that $T_i(\text{Fix}(\phi) \cap \text{EP}(\mathcal{J})) \subset \text{Fix}(\phi)$ for each $i \in \mathbb{N}$ and $\mathcal{F} = \bigcap_{i=1}^{\infty} \text{Fix}(T_i) \cap \text{Fix}(\phi) \cap \text{EP}(\mathcal{J}) \cap \text{VI}(C, A) \cap \text{VI}(C, B) \neq \emptyset$. Let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ and $\{\eta_n\}$ be a sequences in $(0, 1)$. Let $\{\zeta_n\}$ a sequence in $(0, 2\beta)$, $\{\delta_n\}$ a sequence in $(0, 2\gamma)$, $\{r_{k,n}\}_{k=1}^M$ be sequences in $(0, \infty)$ and $\{\lambda_n\}$ a sequence of real numbers such that $0 < \lambda_n \leq b < 1$. Assume that,

- (B₁) $\lim_{n \rightarrow \infty} \eta_n = \eta \in (0, 1)$, $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$,
 (B₂) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$,
 (B₃) $\alpha_n + \beta_n + \gamma_n = 1$,
 (B₄) $\lim_{n \rightarrow \infty} |\zeta_{n+1} - \zeta_n| = \lim_{n \rightarrow \infty} |\delta_{n+1} - \delta_n| = 0$,
 (B₅) $\liminf_{n \rightarrow \infty} r_{k,n} > 0$ and $\lim_{n \rightarrow \infty} (r_{k,n+1} - r_{k,n}) = 0$ for $k \in \{1, 2, \dots, M\}$.

Let f be a contraction of C into itself with coefficient $\alpha \in (0, 1)$ and given $x_1 \in C$ arbitrarily. If the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ are generated iteratively by $x_1 \in C$ and

$$\begin{cases} z_n = J_{T_M, n}^{F_M} \dots J_{T_2, n}^{F_2} J_{T_1, n}^{F_1} x_n, \\ \gamma_n = \eta_n P_C(z_n - \zeta_n A z_n) + (1 - \eta_n) P_C(z_n - \delta_n B z_n), \\ x_{n+1} = \alpha_n f(T_{\mu_n} W_n \gamma_n) + \beta_n x_n + \gamma_n T_{\mu_n} W_n \gamma_n, \quad n \geq 1, \end{cases} \quad (11)$$

then, the sequences $\{x_n\}$, $\{y_n\}$ and $\{J_{T_{k,n}}^{F_k} x_n\}_{k=1}^M$ converge strongly to $x^* \in \mathcal{F}$, which is the unique solution of the system of variational inequalities:

$$\begin{cases} \langle f(x^*) - x^*, x^* - y \rangle \geq 0, \quad \forall y \in \mathcal{F}, \\ \langle Bx^*, y - x^* \rangle \geq 0 \quad \forall y \in C, \\ \langle Ax^*, y - x^* \rangle \geq 0 \quad \forall y \in C. \end{cases}$$

Proof. Since A is a β -inverse strongly monotone map, for any $x, y \in C$, we have

$$\begin{aligned} & \| (I - \zeta_n A)x - (I - \zeta_n A)y \|^2 \\ &= \| (x - y) - \zeta_n (Ax - Ay) \|^2 \\ &= \| x - y \|^2 - 2\zeta_n \langle x - y, Ax - Ay \rangle + \zeta_n^2 \| Ax - Ay \|^2 \\ &\leq \| x - y \|^2 - 2\zeta_n \beta \| Ax - Ay \|^2 + \zeta_n^2 \| Ax - Ay \|^2 \\ &= \| x - y \|^2 + \zeta_n (\zeta_n - 2\beta) \| Ax - Ay \|^2 \\ &\leq \| x - y \|^2 \end{aligned}$$

It follows that

$$\| (I - \zeta_n A)x - (I - \zeta_n A)y \| \leq \| x - y \|. \quad (12)$$

Since B is a β -inverse strongly monotone map, repeating the same argument as above, we can deduce that

$$\| (I - \delta_n B)x - (I - \delta_n B)y \| \leq \| x - y \|. \quad (13)$$

Let $p \in \mathcal{F}$, in the context of the variational inequality problem the characterization of projection (9) implies that $p = P_C(p - \zeta_n A p)$ and $p = P_C(p - \delta_n B p)$. Using (12) and (13), we get

$$\begin{aligned} \| y_n - p \| &= \| \eta_n P_C(z_n - \zeta_n A z_n) + (1 - \eta_n) P_C(z_n - \delta_n B z_n) - p \| \\ &= \| \eta_n [P_C(z_n - \zeta_n A z_n) - P_C(p - \zeta_n A p)] \\ &\quad + (1 - \eta_n) [P_C(z_n - \delta_n B z_n) - P_C(p - \delta_n B p)] \| \\ &\leq \eta_n \| P_C(z_n - \zeta_n A z_n) - P_C(p - \zeta_n A p) \| \\ &\quad + (1 - \eta_n) \| P_C(z_n - \delta_n B z_n) - P_C(p - \delta_n B p) \| \\ &\leq \eta_n \| z_n - p \| + (1 - \eta_n) \| z_n - p \| = \| z_n - p \|. \end{aligned} \quad (14)$$

By taking $v_n = P_C(z_n - \zeta_n A z_n)$, $w_n = P_C(z_n - \delta_n B z_n)$ and $\mathcal{J}_n^k = J_{r_{k,n}}^{F_k} \cdots J_{r_{2,n}}^{F_2} J_{r_{1,n}}^{F_1}$ for $k \in \{1, 2, \dots, M\}$ and $\mathcal{J}_n^0 = I$ for all $n \in \mathbb{N}$, we shall equivalently write scheme (11) as follows:

$$\begin{aligned} z_n &= \mathcal{J}_n^M x_n, \\ \gamma_n &= \eta_n v_n + (1 - \eta_n) w_n, \\ x_{n+1} &= \alpha_n f(T_{\mu_n} W_n \gamma_n) + \beta_n x_n + \gamma_n T_{\mu_n} W_n \gamma_n, \quad n \geq 1. \end{aligned}$$

We shall divide the proof into several steps.

Step 1. The sequence $\{x_n\}$ is bounded.

Proof of Step 1. Let $p \in \mathcal{F}$. Since for each $k \in \{1, 2, \dots, M\}$, $J_{r_{k,n}}^{F_k}$ is nonexpansive we have

$$\|\mathcal{J}_n^k x_n - p\| = \|\mathcal{J}_n^k x_n - \mathcal{J}_n^k p\| \leq \|x_n - p\|, \quad \forall k \in \{1, 2, \dots, M\}. \quad (15)$$

Thus, by Lemmas 2.1, 2.5 and (14), we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\| + \beta_n \|x_n - p\| + \gamma_n \|T_{\mu_n} W_n \mathcal{J}_n^M \gamma_n - p\| \\ &\leq \alpha_n [\|f(T_{\mu_n} W_n \gamma_n) - f(p)\| + \|f(p) - p\|] + \beta_n \|x_n - p\| + \gamma_n \|\gamma_n - p\| \\ &\leq \alpha_n \alpha \|x_n - p\| + \alpha_n \|f(p) - p\| + (\beta_n + \gamma_n) \|x_n - p\| \\ &= [1 - \alpha_n(1 - \alpha)] \|x_n - p\| + \alpha_n \|f(p) - p\| \\ &\leq \max \left\{ \|x_n - p\|, \frac{1}{1 - \alpha} \|f(p) - p\| \right\}. \end{aligned}$$

By induction,

$$\|x_n - p\| \leq \max \left\{ \|x_1 - p\|, \frac{1}{1 - \alpha} \|f(p) - p\| \right\}, \quad n \geq 1.$$

Step 2. Let $\{u_n\}$ be a bounded sequence in H . Then

$$\lim_{n \rightarrow \infty} \|\mathcal{J}_{n+1}^k u_n - \mathcal{J}_n^k u_n\| = 0, \quad (16)$$

for every $k \in \{1, 2, \dots, M\}$.

Proof of Step 2. This assertion is proved in [27, Step 2].

Step 3. Let $\{u_n\}$ be a bounded sequence in H . Then

$$\lim_{n \rightarrow \infty} \|W_{n+1} u_n - W_n u_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|T_{\mu_{n+1}} u_n - T_{\mu_n} u_n\| = 0.$$

This assertion is proved in [21, Step 3].

Step 4. $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$.

Proof of Step 4. Setting $x_{n+1} = \beta_n x_n + (1 - \beta_n) t_n$ for all $n \geq 1$, we have

$$\begin{aligned} t_{n+1} - t_n &= \frac{1}{1 - \beta_{n+1}} [x_{n+2} - \beta_{n+1} x_{n+1}] - \frac{1}{1 - \beta_n} [x_{n+1} - \beta_n x_n] \\ &= \frac{\alpha_{n+1}}{1 - \beta_{n+1}} [f(T_{\mu_{n+1}} W_{n+1} \gamma_{n+1}) - f(T_{\mu_n} W_n \gamma_n)] + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) f(T_{\mu_n} W_n \gamma_n) \\ &\quad + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} [T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_n} W_n \gamma_n] \\ &\quad + \left[\frac{\gamma_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_n}{1 - \beta_n} \right] T_{\mu_n} W_n \gamma_n. \end{aligned}$$

Therefore, we have

$$\begin{aligned} & \|t_{n+1} - t_n\| \\ & \leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}} [\alpha \|T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_n} W_n \gamma_n\| + \|T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_n} W_n \gamma_n\|] \\ & \quad + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| [\|f(T_{\mu_n} W_n \gamma_n)\| + \|T_{\mu_n} W_n \gamma_n\|] \\ & \quad + \|T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_n} W_n \gamma_n\|. \end{aligned}$$

On the other hand

$$\begin{aligned} & \|T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_n} W_n \gamma_n\| \\ & \leq \|T_{\mu_{n+1}} W_{n+1} \gamma_{n+1} - T_{\mu_{n+1}} W_{n+1} \gamma_n\| \\ & \quad + \|T_{\mu_{n+1}} W_{n+1} \gamma_n - T_{\mu_{n+1}} W_n \gamma_n\| \\ & \quad + \|T_{\mu_{n+1}} W_n \gamma_n - T_{\mu_n} W_n \gamma_n\| \\ & \leq \|\gamma_{n+1} - \gamma_n\| + \|W_{n+1} \gamma_n - W_n \gamma_n\| + \|T_{\mu_{n+1}} W_n \gamma_n - T_{\mu_n} W_n \gamma_n\| \end{aligned}$$

Observing that $z_n = \mathcal{J}_n^M x_n$, $z_{n+1} = \mathcal{J}_{n+1}^M x_{n+1}$ and $\mathcal{J}_n^M x_n = J_{r_{M,n}}^{F_M} \mathcal{J}_n^{M-1} x_n$ we get

$$\frac{1}{r_{M,n}} \langle \gamma - z_n, z_n - \mathcal{J}_n^{M-1} x_n \rangle + F_M(z_n, \gamma) \geq 0, \quad \forall \gamma \in C, \quad (17)$$

and

$$\frac{1}{r_{M,n+1}} \langle \gamma - z_{n+1}, z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1} \rangle + F_M(z_{n+1}, \gamma) \geq 0, \quad \forall \gamma \in C, \quad (18)$$

Take $\gamma = z_{n+1}$ in (17) and $\gamma = z_n$ in (18), by using (A_2) , it follows that

$$\left\langle z_{n+1} - z_n, z_n - \mathcal{J}_n^{M-1} x_n - \frac{r_{M,n}}{r_{M,n+1}} (z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1}) \right\rangle \geq 0,$$

and hence

$$\begin{aligned} & \langle z_{n+1} - z_n, z_n - \mathcal{J}_n^{M-1} x_n - z_{n+1} + \mathcal{J}_{n+1}^{M-1} x_{n+1} \\ & \quad + \left(1 - \frac{r_{M,n}}{r_{M,n+1}} \right) (z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1}) \rangle \geq 0, \end{aligned}$$

Thus, we have

$$\begin{aligned} & \|z_{n+1} - z_n\| \\ & \leq \|\mathcal{J}_{n+1}^{M-1} x_{n+1} - \mathcal{J}_n^{M-1} x_n\| + \left| 1 - \frac{r_{M,n}}{r_{M,n+1}} \right| \|z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1}\| \\ & \leq \|\mathcal{J}_{n+1}^{M-1} x_{n+1} - \mathcal{J}_{n+1}^{M-1} x_n\| + \|\mathcal{J}_{n+1}^{M-1} x_n - \mathcal{J}_n^{M-1} x_n\| \\ & \quad + \left| 1 - \frac{r_{M,n}}{r_{M,n+1}} \right| \|z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1}\| \\ & \leq \|x_{n+1} - x_n\| + \|\mathcal{J}_{n+1}^{M-1} x_n - \mathcal{J}_n^{M-1} x_n\| \\ & \quad + \left| 1 - \frac{r_{M,n}}{r_{M,n+1}} \right| \|z_{n+1} - \mathcal{J}_{n+1}^{M-1} x_{n+1}\|. \end{aligned}$$

Since $v_n = P_C(z_n - \zeta_n A z_n)$ and $w_n = P_C(z_n - \delta_n B z_n)$, it follows from the definition of $\{y_n\}$ that

$$\begin{aligned}
& \|y_{n+1} - y_n\| \\
&= \|\eta_{n+1}v_{n+1} + (1 - \eta_{n+1})w_{n+1} - \eta_nv_n - (1 - \eta_n)w_n\| \\
&= \|\eta_{n+1}(v_{n+1} - v_n) + (\eta_{n+1} - \eta_n)v_n + (1 - \eta_{n+1})w_{n+1} \\
&\quad - (1 - \eta_{n+1})w_n + (\eta_n - \eta_{n+1})w_n\| \\
&\leq \eta_{n+1}\|v_{n+1} - v_n\| + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) \\
&\quad + (1 - \eta_{n+1})\|w_{n+1} - w_n\| \\
&= \eta_{n+1}\|P_C(z_{n+1} - \zeta_{n+1}Az_{n+1}) - P_C(z_n - \zeta_nAz_n)\| \\
&\quad + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) \\
&\quad + (1 - \eta_{n+1})\|P_C(z_{n+1} - \delta_{n+1}Bz_{n+1}) - P_C(z_n - \delta_nBz_n)\| \\
&= \eta_{n+1}\|P_C(z_{n+1} - \zeta_{n+1}Az_{n+1}) - P_C(z_n - \zeta_nAz_n) \\
&\quad + P_C(z_n - \zeta_{n+1}Az_n) - P_C(z_n - \zeta_nAz_n)\| \\
&\quad + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) \\
&\quad + (1 - \eta_{n+1})\|P_C(z_{n+1} - \delta_{n+1}Bz_{n+1}) - P_C(z_n - \delta_nBz_n) \\
&\quad + P_C(z_n - \delta_{n+1}Bz_n) - P_C(z_n - \delta_nBz_n)\| \\
&\leq \eta_{n+1}\|z_{n+1} - z_n\| + \eta_{n+1}|\zeta_{n+1} - \zeta_n|\|Az_n\| \\
&\quad + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) + (1 - \eta_{n+1})\|z_{n+1} - z_n\| \\
&\quad + (1 - \eta_{n+1})|\delta_{n+1} - \delta_n|\|Bz_n\| \\
&\leq \|z_{n+1} - z_n\| + \eta_{n+1}|\zeta_{n+1} - \zeta_n|\|Az_n\| \\
&\quad + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) + |\delta_{n+1} - \delta_n|\|Bz_n\|.
\end{aligned}$$

Therefore,

$$\begin{aligned}
& \|t_{n+1} - t_n\| - \|x_{n+1} - x_n\| \\
&\leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}}[\|y_{n+1} - y_n\| + \|T_{\mu_{n+1}}W_{n+1}y_{n+1} - T_{\mu_n}W_ny_n\|] \\
&\quad + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| [\|f(T_{\mu_n}W_ny_n)\| + \|T_{\mu_n}W_ny_n\|] \\
&\quad + \|\mathcal{J}_{n+1}^{M-1}x_n - \mathcal{J}_n^{M-1}x_n\| + \left| 1 - \frac{r_{M,n}}{r_{M,n+1}} \right| \|z_{n+1} - \mathcal{J}_{n+1}^{M-1}x_{n+1}\| \\
&\quad + \eta_{n+1}|\zeta_{n+1} - \zeta_n|\|Az_n\| + |\eta_{n+1} - \eta_n|(\|v_n\| + \|w_n\|) \\
&\quad + |\delta_{n+1} - \delta_n|\|Bz_n\| + \|W_{n+1}y_n - W_ny_n\| + \|T_{\mu_{n+1}}W_ny_n - T_{\mu_n}W_ny_n\|.
\end{aligned}$$

This together with conditions (B_1) , (B_4) , Steps 2 and 3 imply that

$$\limsup_{n \rightarrow \infty} (\|t_{n+1} - t_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Hence by Lemma 2.4, we obtain $\lim_{n \rightarrow \infty} \|t_n - x_n\| = 0$. Consequently,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = (1 - \beta_n)\|t_n - x_n\| = 0.$$

Step 5. $\lim_{n \rightarrow \infty} \|\mathcal{J}_n^{k+1}x_n - \mathcal{J}_n^kx_n\| = 0, \forall k \in \{0, 1, 2, \dots, M-1\}$.

Proof of Step 5. Let $p \in \mathcal{F}$ and $k \in \{1, 2, \dots, M-1\}$. Since $J_{\Gamma_{k+1,n}}^{F_{k+1}}$ is firmly nonexpansive, we obtain

$$\begin{aligned}
& \|\mathcal{J}_n^{k+1}x_n - p\|^2 \\
&= \|J_{\Gamma_{k+1,n}}^{F_{k+1}}\mathcal{J}_n^kx_n - J_{\Gamma_{k+1,n}}^{F_{k+1}}p\|^2 = \langle J_{\Gamma_{k+1,n}}^{F_{k+1}}\mathcal{J}_n^kx_n - p, \mathcal{J}_n^kx_n - p \rangle \\
&= \frac{1}{2} \left[\|J_{\Gamma_{k+1,n}}^{F_{k+1}}\mathcal{J}_n^kx_n - p\|^2 + \|\mathcal{J}_n^kx_n - p\|^2 - \|J_{\Gamma_{k+1,n}}^{F_{k+1}}\mathcal{J}_n^kx_n - \mathcal{J}_n^kx_n\|^2 \right].
\end{aligned}$$

It follows that

$$\|\mathcal{J}_n^{k+1}x_n - p\|^2 \leq \|x_n - p\|^2 - \|\mathcal{J}_n^{k+1}x_n - \mathcal{J}_n^kx_n\|^2. \quad (19)$$

Using Lemma 2.9, (14) and (19), we obtain

$$\begin{aligned} & \|x_{n+1} - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|T_{\mu_n}W_n\gamma_n - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\gamma_n - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ & = \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\mathcal{J}_n^Mx_n - p\|^2 \\ & = \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n \|J_{r_{M,n}}^{F_M} \cdots J_{r_{k+2,n}}^{F_{k+2}} \mathcal{J}_n^{k+1}\gamma_n - J_{r_{M,n}}^{F_M} \cdots J_{r_{k+2,n}}^{F_{k+2}} p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\mathcal{J}_n^{k+1}\gamma_n - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\mathcal{J}_n^{k+1}x_n - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n [\|x_n - p\|^2 - \|\mathcal{J}_n^{k+1}x_n - \mathcal{J}_n^kx_n\|]. \end{aligned}$$

Then, we have

$$\begin{aligned} & \gamma_n \|\mathcal{J}_n^{k+1}x_n - \mathcal{J}_n^kx_n\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + (1 - \alpha_n - \beta_n) \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \\ & = \alpha_n [\|f(T_{\mu_n}W_n\gamma_n) - p\|^2 - \|x_n - p\|^2] + \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \\ & \leq \alpha_n [\|f(T_{\mu_n}W_n\gamma_n) - p\|^2 - \|x_n - p\|^2] \\ & \quad + \|x_n - x_{n+1}\| [\|x_n - p\| + \|x_{n+1} - p\|]. \end{aligned}$$

It is easily seen that $\liminf_{n \rightarrow \infty} \gamma_n > 0$. So we have

$$\lim_{n \rightarrow \infty} \|\mathcal{J}_n^{k+1}x_n - \mathcal{J}_n^kx_n\| = 0.$$

Step 6. $\lim_{n \rightarrow \infty} \|x_n - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\| = 0$.

Proof of Step 6. Observe that

$$\begin{aligned} & \|x_n - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\| \\ & \leq \|x_n - x_{n+1}\| + \|x_{n+1} - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\| \\ & = \|x_n - x_{n+1}\| + \|\alpha_n [f(T_{\mu_n}W_n\gamma_n) - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n] \\ & \quad + \beta_n [x_n - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n]\| \\ & \leq \|x_n - x_{n+1}\| + \alpha_n [\|f(T_{\mu_n}W_n\gamma_n)\| + \|T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\|] \\ & \quad + \beta_n \|x_n - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\|, \end{aligned}$$

hence

$$\begin{aligned} & \|x_n - T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\| \\ & \leq \frac{1}{1 - \beta_n} \|x_{n+1} - x_n\| + \frac{\alpha_n}{1 - \beta_n} [\|f(T_{\mu_n}W_n\gamma_n)\| + \|T_{\mu_n}W_n\mathcal{J}_n^M\gamma_n\|]. \end{aligned}$$

It follows from conditions (B_1) , (B_2) and Step 4, that

$$\lim_{n \rightarrow \infty} \|x_n - T_{\mu_n} W_n \mathcal{J}_n^M y_n\| = 0.$$

Step 7. $\lim_{n \rightarrow \infty} \|x_n - T_t x_n\| = 0$, for all $t \in S$.

Proof of Step 7. Let $p \in \mathcal{F}$ and set $M_0 = \max\{\|x_1 - p\|, \frac{1}{1-\alpha} \|f(p) - p\|\}$ and $D = \{y \in H : \|y - p\| \leq M_0\}$, we remark that D is bounded closed convex set, $\{y_n\} \subset D$ and it is invariant under $\{J_{t_{k,n}}^{F_k} : k = 1, 2, \dots, M, \forall n \in \mathbb{N}\}$, ϕ and W_n for all $n \in \mathbb{N}$. We will show that

$$\limsup_{n \rightarrow \infty} \sup_{y \in D} \|T_{\mu_n} y - T_t T_{\mu_n} y\| = 0, \quad \forall t \in S \quad (20)$$

Let $\epsilon > 0$. By [33, Theorem 1.2], there exists $\delta > 0$ such that

$$\overline{\text{co}}F_\delta(T_t; D) + B_\delta \subset F_\epsilon(T_t; D), \quad \forall t \in S. \quad (21)$$

Also by [33, Corollary 1.1], there exists a natural number N such that

$$\left\| \frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y - T_t \left(\frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y \right) \right\| \leq \delta, \quad (22)$$

for all $t, s \in S$ and $y \in D$. Let $t \in S$. Since $\{\mu_n\}$ is strongly left regular, there exists $N_0 \in \mathbb{N}$ such that $\|\mu_n - I_{t^i}^* \mu_n\| \leq \frac{\delta}{(M_0 + \|p\|)}$ for $n \geq N_0$ and $i = 1, 2, \dots, N$. Then, we have

$$\begin{aligned} & \sup_{y \in D} \left\| T_{\mu_n} y - \int \frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y d\mu_n(s) \right\| \\ &= \sup_{y \in D} \sup_{\|z\|=1} \left| \langle T_{\mu_n} y, z \rangle - \left\langle \int \frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y d\mu_n(s), z \right\rangle \right| \\ &= \sup_{y \in D} \sup_{\|z\|=1} \left| \frac{1}{N+1} \sum_{i=0}^N (\mu_n)_s \langle T_s y, z \rangle - \frac{1}{N+1} \sum_{i=0}^N (\mu_n)_s \langle T_{t^i s} y, z \rangle \right| \\ &\leq \frac{1}{N+1} \sum_{i=0}^N \sup_{y \in D} \sup_{\|z\|=1} |(\mu_n)_s \langle T_s y, z \rangle - (I_{t^i}^* \mu_n)_s \langle T_s y, z \rangle| \\ &\leq \max_{i=1,2,\dots,N} \|\mu_n - I_{t^i}^* \mu_n\| (M_0 + \|p\|) \leq \delta, \quad \forall n \geq N_0. \end{aligned} \quad (23)$$

By Lemma 2.1 we have

$$\int \frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y d\mu_n(s) \in \overline{\text{co}} \left\{ \frac{1}{N+1} \sum_{i=0}^N T_{t^i} (T_s y) : s \in S \right\}. \quad (24)$$

It follows from (21), (22), (23) and (24) that

$$\begin{aligned} T_{\mu_n} y &\in \overline{\text{co}} \left\{ \frac{1}{N+1} \sum_{i=0}^N T_{t^i s} y : s \in S \right\} + B_\delta \\ &\subset \overline{\text{co}}F_\delta(T_t; D) + B_\delta \subset F_\epsilon(T_t; D), \end{aligned}$$

for all $y \in D$ and $n \geq N_0$. Therefore,

$$\limsup_{n \rightarrow \infty} \sup_{y \in D} \|T_t(T_{\mu_n}y) - T_{\mu_n}y\| \leq \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, we get (20).

Let $t \in S$ and $\varepsilon > 0$. Then, there exists $\delta > 0$, which satisfies (21). From condition (B_1) , (20) and Step 6, there exists $N_1 \in \mathbb{N}$ such that $\alpha_n < \frac{\delta}{4M_0}$, $T_{\mu_n}y \in F_\delta(T_t, D)$ for all $y \in D$ and $\|x_n - T_{\mu_n}W_n y_n\| < \frac{\delta}{2}$ for all $n \geq N_1$. We note that

$$\begin{aligned} \alpha_n \|f(T_{\mu_n}W_n y_n) - T_{\mu_n}W_n y_n\| \\ \leq \alpha_n [\|f(T_{\mu_n}W_n y_n) - f(p)\| + \|f(p) - p\| + \|p - T_{\mu_n}W_n y_n\|] \\ \leq \alpha_n [\alpha \|y_n - p\| + \|f(p) - p\| + \|p - y_n\|] \leq 2M_0 \alpha_n \leq \frac{\delta}{2}, \end{aligned}$$

for all $n \geq N_1$. Therefore, we have

$$\begin{aligned} x_{n+1} &= T_{\mu_n}W_n y_n + \alpha_n (f(T_{\mu_n}W_n y_n) - T_{\mu_n}W_n y_n) \\ &\quad + \beta_n (x_n - T_{\mu_n}W_n y_n) \\ &\in F_\delta(T_t; D) + B_{\frac{\delta}{2}} + B_{\frac{\delta}{2}} \subset F_\delta(T_t; D) + B_\delta \subset F_\varepsilon(T_t; D), \end{aligned}$$

for all $n \geq N_1$. This shows that

$$\|x_n - T_t x_n\| \leq \varepsilon, \quad \forall n \geq N_1.$$

Since $\varepsilon > 0$ is arbitrary, we get $\lim_{n \rightarrow \infty} \|x_n - T_t(x_n)\| = 0$.

Step 8. The weak ω -limit set of $\{x_n\}$, $\omega_\omega\{x_n\}$, is a subset of \mathcal{F} .

Proof of Step 8. Let $z \in \omega_\omega\{x_n\}$ and let $\{x_{n_m}\}$ be a subsequence of $\{x_n\}$ weakly converging to z , we need to show that $z \in \mathcal{F}$. Noting Step 5, with no loss of generality, we may assume that $\mathcal{J}_{n_m}^k x_{n_m} \rightharpoonup z, \forall k \in \{1, 2, \dots, M\}$. At first, note that by (A_2) and given $y \in C$ and $k \in \{1, 2, \dots, M\}$, we have

$$\left\langle y - \mathcal{J}_{n_m}^{k+1} x_{n_m}, \frac{1}{r_{k+1, n_m}} (\mathcal{J}_{n_m}^{k+1} x_{n_m} - \mathcal{J}_{n_m}^k x_{n_m}) \right\rangle \geq F_{k+1}(y, \mathcal{J}_{n_m}^{k+1} x_{n_m}).$$

Step 5 and condition (B_5) imply that

$$\frac{\mathcal{J}_{n_m}^{k+1} x_{n_m} - \mathcal{J}_{n_m}^k x_{n_m}}{r_{k+1, n_m}} \rightarrow 0.$$

Since $\mathcal{J}_{n_m}^k x_{n_m} \rightharpoonup z$, from the lower semi-continuity of F_{k+1} on the second variable, we have $F_{k+1}(y, z) \leq 0$ for all $y \in C$ and for all $k \in \{0, 1, 2, \dots, M-1\}$. For t with $0 < t \leq 1$ and $y \in C$, let $y_t = ty + (1-t)z$. Since $y \in C$ and $z \in C$, we have $y_t \in C$ and hence $F_{k+1}(y_t, z) \leq 0$. So from the convexity of F_{k+1} on second variable, we have

$$0 = F_{k+1}(y_t, y_t) \leq tF_{k+1}(y_t, y) + (1-t)F_{k+1}(y_t, z) \leq tF_{k+1}(y_t, y) \leq F_{k+1}(y_t, y).$$

hence $F_{k+1}(y_t, y) \geq 0$. therefore, we have $F_{k+1}(z, y) \geq 0$ for all $y \in C$ and $k \in \{0, 1, 2, \dots, M-1\}$. Therefore $z \in \bigcap_{k=1}^M \text{EP}(F_k) = \text{EP}(\mathcal{J})$.

Since $x_{n_m} \rightharpoonup z$, it follows by Step 7 and Lemma 2.2 that $z \in \text{Fix}(T_t)$ for all $t \in S$. Therefore, $z \in \text{Fix}(\phi)$. We will show $z \in \text{Fix}(W)$. Assume $z \notin \text{Fix}(W)$ Since

$z \in \text{Fix}(\varphi) \cap \text{EP}(\mathcal{J})$, by our assumption, we have $T_i z \in \text{Fix}(\phi), \forall i \in \mathbb{N}$ and then $W_n z \in \text{Fix}(\phi)$. Hence by Lemma 2.1, $T_{\mu_n} W_n z = W_n z$, therefore by Lemma 2.5, we get

$$T_{\mu_n} W_n \mathcal{J}_n^M z = W_n z, \quad \forall n \in \mathbb{N}. \quad (25)$$

Now, by (25), Step 6, Lemma 2.6 and Opial's condition, we have

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \|x_{n_m} - z\| \\ & < \liminf_{n \rightarrow \infty} \|x_{n_m} - Wz\| \\ & \leq \liminf_{n \rightarrow \infty} (\|x_{n_m} - T_{\mu_{n_m}} W_{n_m} \mathcal{J}_{n_m}^M x_{n_m}\| \\ & \quad + \|T_{\mu_{n_m}} W_{n_m} \mathcal{J}_{n_m}^M x_{n_m} - T_{\mu_{n_m}} W_{n_m} \mathcal{J}_{n_m}^M z\| + \|T_{\mu_{n_m}} W_{n_m} \mathcal{J}_{n_m}^M z - Wz\|) \\ & \leq \liminf_{n \rightarrow \infty} (\|x_{n_m} - T_{\mu_{n_m}} W_{n_m} \mathcal{J}_{n_m}^M x_{n_m}\| + \|x_{n_m} - z\| + \|W_{n_m} z - Wz\|) \\ & \leq \liminf_{n \rightarrow \infty} \|x_{n_m} - z\|. \end{aligned}$$

This is a contradiction. So we get $z \in \text{Fix}(W) = \bigcap_{i=1}^{\infty} \text{Fix}(T_i)$.

Now, let us show that $z \in \text{VI}(C, A) \cap \text{VI}(C, B)$. Observe that,

$$\begin{aligned} & \|x_{n+1} - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|T_{\mu_n} W_n \mathcal{J}_n^M \gamma_n - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\gamma_n - p\|^2 \\ & = \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\eta_n P_C(z_n - \zeta_n A z_n) \\ & \quad + (1 - \eta_n) P_C(z_n - \delta_n B z_n) - p\|^2 \\ & = \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|\eta_n [P_C(z_n - \zeta_n A z_n) \\ & \quad - P_C(p - \zeta_n A p)] + (1 - \eta_n) [P_C(z_n - \delta_n B z_n) - P_C(p - \delta_n B p)]\|^2. \end{aligned} \quad (26)$$

From (26), we have

$$\begin{aligned} & \|x_{n+1} - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n [\eta_n \|(z_n - p) - \zeta_n (A z_n - A p)\|^2 + (1 - \eta_n) \|z_n - p\|^2] \\ & = \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n (1 - \eta_n) \|z_n - p\|^2 + \gamma_n \eta_n [\|z_n - p\|^2 + \zeta_n^2 \|A z_n - A p\|^2 \\ & \quad - 2 \zeta_n \langle A z_n - A p, z_n - p \rangle] \\ & \leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n (1 - \eta_n) \|z_n - p\|^2 + \gamma_n \eta_n [\|z_n - p\|^2 + \zeta_n^2 \|A z_n - A p\|^2 \\ & \quad - 2 \zeta_n \beta \|A z_n - A p, z_n - p\|^2] \\ & = \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 \\ & \quad + \gamma_n \|z_n - p\|^2 + \zeta_n (\zeta_n - 2\beta) \|A z_n - A p\|^2 \\ & \leq \|x_n - p\|^2 + \alpha_n [\|f(T_{\mu_n} W_n \gamma_n) - p\|^2 - \|x_n - p\|^2] \\ & \quad + \zeta_n (\zeta_n - 2\beta) \|A z_n - A p\|^2, \end{aligned}$$

which implies that

$$\begin{aligned} & -\zeta_n(\zeta_n - 2\beta)\|Az_n - Ap\|^2 \\ & \leq [\|x_n - p\| + \|x_{n+1} - p\|]\|x_n - x_{n+1}\| \\ & \quad + \alpha_n[\|f(T_{\mu_n}W_n\gamma_n) - p\|^2 - \|x_n - p\|^2]. \end{aligned}$$

Therefore, from step 4 and condition B_1 , we obtain

$$\lim_{n \rightarrow \infty} \|Az_n - Ap\| = 0. \quad (27)$$

On the other hand from (26), we have

$$\begin{aligned} & \|x_{n+1} - p\|^2 \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n [\eta_n \|z_n - p\|^2 \\ & \quad + (1 - \eta_n) \|(z_n - p) - \delta_n(Bz_n - Bp)\|^2] \\ & = \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n [\eta_n \|z_n - p\|^2 \\ & \quad + (1 - \eta_n)(\|z_n - p\|^2 - 2\delta_n \langle Bz_n - Bp, z_n - p \rangle + \delta_n^2 \|Bz_n - Bp\|^2)] \\ & \leq \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n [\eta_n \|z_n - p\|^2 \\ & \quad + (1 - \eta_n)(\|z_n - p\|^2 - 2\delta_n \gamma \|Bz_n - Bp\|^2 + \delta_n^2 \|Bz_n - Bp\|^2)] \\ & = \alpha_n \|f(T_{\mu_n}W_n\gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ & \quad + \delta_n(\delta_n - 2\gamma)\gamma_n(1 - \eta_n) \|Bz_n - Bp\|^2 \\ & \leq \|x_n - p\|^2 + \alpha_n[\|f(T_{\mu_n}W_n\gamma_n) - p\|^2 - \|x_n - p\|^2] \\ & \quad + \delta_n(\delta_n - 2\gamma) \|Bz_n - Bp\|^2 \end{aligned}$$

which implies that

$$\begin{aligned} & -\delta_n(\delta_n - 2\gamma) \|Bz_n - Bp\|^2 \\ & \leq [\|x_n - p\| + \|x_{n+1} - p\|]\|x_n - x_{n+1}\| \\ & \quad + \alpha_n[\|f(T_{\mu_n}W_n\gamma_n) - p\|^2 - \|x_n - p\|^2]. \end{aligned}$$

Therefore, from step 4 and condition B_1 , we obtain

$$\lim_{n \rightarrow \infty} \|Bz_n - Bp\| = 0. \quad (28)$$

From (6) and (12), we have

$$\begin{aligned} & \|v_n - p\|^2 \\ & = \|P_C(z_n - \zeta_n Az_n) - P_C(p - \zeta_n Ap)\|^2 \\ & \leq \langle (z_n - \zeta_n Az_n) - (p - \zeta_n Ap), v_n - p \rangle \\ & = \frac{1}{2} [\|(z_n - \zeta_n Az_n) - (p - \zeta_n Ap)\|^2 + \|v_n - p\|^2 \\ & \quad - \|(z_n - \zeta_n Az_n) - (p - \zeta_n Ap) - (v_n - p)\|^2] \\ & = \frac{1}{2} [\|z_n - p\|^2 + \|v_n - p\|^2 - \|z_n - v_n\|^2 \\ & \quad + 2\zeta_n \langle z_n - v_n, Az_n - Ap \rangle - \zeta_n^2 \|Az_n - Ap\|^2]. \end{aligned}$$

So we obtain

$$\begin{aligned}\|v_n - p\|^2 &\leq \|z_n - p\|^2 - \|z_n - v_n\|^2 \\ &\quad + 2\zeta_n \langle z_n - v_n, Az_n - Ap \rangle - \zeta_n^2 \|Az_n - Ap\|^2.\end{aligned}\quad (29)$$

By using the same method as (29), we have

$$\begin{aligned}\|w_n - p\|^2 &\leq \|z_n - p\|^2 - \|z_n - w_n\|^2 \\ &\quad + 2\delta_n \langle z_n - w_n, Bz_n - Bp \rangle - \delta_n^2 \|Bz_n - Bp\|^2.\end{aligned}\quad (30)$$

From (29), (30) and definition of y_n , we have,

$$\begin{aligned}\|y_n - p\|^2 &= \|\eta_n [P_C(z_n - \zeta_n Az_n) - p] \\ &\quad + (1 - \eta_n) [P_C(z_n - \delta_n Bz_n) - p]\|^2 \\ &= \|\eta_n (v_n - p) + (1 - \eta_n) (w_n - p)\|^2 \\ &\leq \eta_n \|v_n - p\|^2 + (1 - \eta_n) \|w_n - p\|^2 \\ &\leq \eta_n [\|z_n - p\|^2 - \|z_n - v_n\|^2 + 2\zeta_n \langle z_n - v_n, Az_n - Ap \rangle \\ &\quad - \zeta_n^2 \|Az_n - Ap\|^2] + (1 - \eta_n) [\|z_n - p\|^2 - \|z_n - w_n\|^2 \\ &\quad + 2\delta_n \langle z_n - w_n, Bz_n - Bp \rangle - \delta_n^2 \|Bz_n - Bp\|^2] \\ &\leq \|z_n - p\|^2 + \eta_n [-\|z_n - v_n\|^2 + 2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| \\ &\quad - \zeta_n^2 \|Az_n - Ap\|^2] + (1 - \eta_n) [-\|z_n - w_n\|^2 \\ &\quad + 2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2]\end{aligned}\quad (31)$$

By (31), we have

$$\begin{aligned}\|x_{n+1} - p\|^2 &\leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|T_{\mu_n} W_n \mathcal{J}_n^M \gamma_n - p\|^2 \\ &\leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|z_n - p\|^2 \\ &\quad + \gamma_n \eta_n [-\|z_n - v_n\|^2 + 2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| \\ &\quad - \zeta_n^2 \|Az_n - Ap\|^2] + \gamma_n (1 - \eta_n) [-\|z_n - w_n\|^2 \\ &\quad + 2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2] \\ &\leq \alpha_n \|f(T_{\mu_n} W_n \gamma_n) - p\|^2 + \beta_n \|x_n - p\|^2 + \gamma_n \|x_n - p\|^2 \\ &\quad - \gamma_n \eta_n \|z_n - v_n\|^2 + \gamma_n \eta_n [2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| \\ &\quad - \zeta_n^2 \|Az_n - Ap\|^2] - \gamma_n (1 - \eta_n) \|z_n - w_n\|^2 \\ &\quad + \gamma_n (1 - \eta_n) [2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2] \\ &= \|x_n - p\|^2 + \alpha_n [\|f(T_{\mu_n} W_n \gamma_n) - p\|^2 - \|x_n - p\|^2] \\ &\quad - \gamma_n \eta_n \|z_n - v_n\|^2 + \gamma_n \eta_n [2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| \\ &\quad - \zeta_n^2 \|Az_n - Ap\|^2] - \gamma_n (1 - \eta_n) \|z_n - w_n\|^2 \\ &\quad + \gamma_n (1 - \eta_n) [2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2],\end{aligned}$$

which implies that

$$\begin{aligned} & \gamma_n \eta_n \|z_n - v_n\|^2 \\ & \leq [\|x_n - p\| + \|x_{n+1} - p\|] \|x_{n+1} - x_n\| \\ & \quad + \alpha_n [\|f(T_{\mu_n} W_n \gamma_n) - p\|^2 - \|x_n - p\|^2] \\ & \quad + \gamma_n \eta_n [2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| - \zeta_n^2 \|Az_n - Ap\|^2] \\ & \quad + \gamma_n (1 - \eta_n) [2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2], \end{aligned}$$

and

$$\begin{aligned} & \gamma_n (1 - \eta_n) \|z_n - w_n\|^2 \\ & \leq [\|x_n - p\| + \|x_{n+1} - p\|] \|x_{n+1} - x_n\| \\ & \quad + \gamma_n \eta_n [2\zeta_n \|z_n - v_n\| \|Az_n - Ap\| - \zeta_n^2 \|Az_n - Ap\|^2] \\ & \quad + \gamma_n (1 - \eta_n) [2\delta_n \|z_n - w_n\| \|Bz_n - Bp\| - \delta_n^2 \|Bz_n - Bp\|^2]. \end{aligned}$$

Therefore, from $0 < \liminf_{n \rightarrow \infty} \gamma_n \leq \limsup_{n \rightarrow \infty} \gamma_n < 1$, condition B_1 , step 4, (27) and (28) we get

$$\lim_{n \rightarrow \infty} \|z_n - v_n\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|z_n - w_n\| = 0. \quad (32)$$

Let $U : H \rightarrow 2^H$ be a set-valued mapping is defined by

$$Ux = \begin{cases} Ax + N_C x, & x \in C, \\ \emptyset, & x \notin C, \end{cases}$$

where $N_C x$ is the normal cone to C at $x \in C$. Since A is monotone. Thus U is maximal monotone see [1]. Let $(x, y) \in G(U)$, hence $y - Ax \in N_C x$ and since $v_n = P_C(z_n - \zeta_n Az_n)$ therefore, $\langle x - v_n, y - Ax \rangle \geq 0$. On the other hand from (7), we have

$$\langle x - v_n, v_n - (z_n - \zeta_n Az_n) \rangle \geq 0,$$

i.e.,

$$\left\langle x - v_n, \frac{v_n - z_n}{\zeta_n} + Az_n \right\rangle \geq 0$$

Therefore, we have

$$\begin{aligned} & \langle x - v_{n_i}, y \rangle \\ & \geq \langle x - v_{n_i}, Ax \rangle \\ & \geq \langle x - v_{n_i}, Ax \rangle - \left\langle x - v_{n_i}, \frac{v_{n_i} - z_{n_i}}{\zeta_{n_i}} + Az_{n_i} \right\rangle \\ & = \left\langle x - v_{n_i}, Ax - \frac{v_{n_i} - z_{n_i}}{\zeta_{n_i}} - Az_{n_i} \right\rangle \\ & = \langle x - v_{n_i}, Ax - Av_{n_i} \rangle + \langle x - v_{n_i}, Av_{n_i} - Az_{n_i} \rangle - \left\langle x - v_{n_i}, \frac{v_{n_i} - z_{n_i}}{\zeta_{n_i}} \right\rangle \\ & \geq \langle x - v_{n_i}, Av_{n_i} - Az_{n_i} \rangle - \left\langle x - v_{n_i}, \frac{v_{n_i} - z_{n_i}}{\zeta_{n_i}} \right\rangle \\ & \geq \langle x - v_{n_i}, Av_{n_i} - Az_{n_i} \rangle - \|x - v_{n_i}\| \left\| \frac{v_{n_i} - z_{n_i}}{\zeta_{n_i}} \right\|. \end{aligned}$$

From (32), we get $\lim_{i \rightarrow \infty} \|v_{n_i} - z_{n_i}\| = 0$. Noting that $x_{n_i} \rightarrow z$ and A is $\frac{1}{\beta}$ -lipschitzian, we obtain

$$\langle x - z, y \rangle \geq 0. \quad (33)$$

Since U is maximal monotone, we have $z \in U^{-1}0$, and hence $z \in \text{VI}(C, A)$. Let $V: H \rightarrow 2^H$ be a set-valued mapping is defined by

$$Vx = \begin{cases} Bx + N_C x, & x \in C, \\ \emptyset, & x \notin C, \end{cases}$$

where $N_C x$ is the normal cone to C at $x \in C$. Since B is monotone. Thus U is maximal monotone see [1]. Repeating the same argument as above, we can derive $z \in \text{VI}(C, B)$. Therefore, $z \in \mathcal{F}$.

Step 9. There exists a unique $x^* \in C$ such that

$$\limsup_{n \rightarrow \infty} \langle f(x^*) - x^*, x_n - x^* \rangle \leq 0.$$

Proof of Step 9. Note that f is a contraction mapping with coefficient $\alpha \in (0, 1)$. Then $\|P_{\mathcal{F}}f(x) - P_{\mathcal{F}}f(y)\| \leq \|f(x) - f(y)\| \leq \alpha\|x - y\|$ for all $x, y \in H$. Therefore $P_{\mathcal{F}}$ is a contraction of H into itself, which implies that there exists a unique element $x^* \in H$ such that $x^* = P_{\mathcal{F}}f(x^*)$. at the same time, we note that $x^* \in C$. Using Lemma 2.3, we have

$$\langle f(x^*) - x^*, x^* - z \rangle \geq 0, \quad \forall z \in \mathcal{F}. \quad (34)$$

We can choose a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle f(x^*) - x^*, x_n - x^* \rangle = \lim_{k \rightarrow \infty} \langle f(x^*) - x^*, x_{n_k} - x^* \rangle.$$

Since $\{x_{n_k}\}$ is bounded, therefore, $\{x_{n_k}\}$ has subsequence $\{x_{n_{k_j}}\}$ such that $x_{n_{k_j}} \rightarrow z^*$. With no loss of generality, we may assume that $x_{n_k} \rightarrow z^*$. Applying Step 8 and (34), we have

$$\limsup_{n \rightarrow \infty} \langle f(x^*) - x^*, x_n - x^* \rangle = \langle f(x^*) - x^*, z^* - x^* \rangle \leq 0.$$

Step 10, The sequences $\{x_n\}$ converges strongly to x^* , which is obtained in Steep 9.

Proof of Step 10. We have

$$\begin{aligned} & \|x_{n+1} - x^*\|^2 \\ &= \|\alpha_n(f(T_{\mu_n}W_n\gamma_n) - x^*) + \beta_n(x_n - x^*) + \gamma_n(T_{\mu_n}W_n\gamma_n - x^*)\|^2 \\ &\leq \|\beta_n(x_n - x^*) + \gamma_n(T_{\mu_n}W_n\gamma_n - x^*)\|^2 + 2\alpha_n\langle f(T_{\mu_n}W_n\gamma_n) - x^*, x_{n+1} - x^* \rangle \\ &\leq [\beta_n\|x_n - x^*\| + \gamma_n\|T_{\mu_n}W_n\gamma_n - x^*\|]^2 \\ &\quad + 2\alpha_n\langle f(T_{\mu_n}W_n\gamma_n) - f(x^*), x_{n+1} - x^* \rangle + 2\alpha_n\langle f(x^*) - x^*, x_{n+1} - x^* \rangle \\ &\leq [\beta_n\|x_n - x^*\| + \gamma_n\|\gamma_n - x^*\|]^2 \\ &\quad + 2\alpha_n\alpha\|\gamma_n - x^*\|\|x_{n+1} - x^*\| + 2\alpha_n\langle f(x^*) - x^*, x_{n+1} - x^* \rangle \\ &\leq [\beta_n\|x_n - x^*\| + \gamma_n\|x_n - x^*\|]^2 \\ &\quad + 2\alpha_n\alpha\|x_n - x^*\|\|x_{n+1} - x^*\| + 2\alpha_n\langle f(x^*) - x^*, x_{n+1} - x^* \rangle \\ &= (1 - \alpha_n)^2\|x_n - x^*\|^2 + \alpha_n\alpha[\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2] \\ &\quad + 2\alpha_n\langle f(x^*) - x^*, x_{n+1} - x^* \rangle \end{aligned}$$

Which implies that

$$\begin{aligned} & \|x_{n+1} - x^*\|^2 \\ & \leq \frac{(1 - \alpha_n)^2 + \alpha_n \alpha}{1 - \alpha_n \alpha} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 - \alpha_n \alpha} \langle f(x^*) - x^*, x_{n+1} - x^* \rangle \\ & = \left(1 - \alpha_n \frac{2 - 2\alpha}{1 - \alpha_n \alpha}\right) \|x_n - x^*\|^2 + \alpha_n \tau_n, \end{aligned} \quad (35)$$

where

$$\tau_n = \frac{\alpha_n}{1 - \alpha_n \alpha} \|x_n - x^*\|^2 + \frac{2}{1 - \alpha_n \alpha} \langle f(x^*) - x^*, x_{n+1} - x^* \rangle,$$

By Step 9, and condition (B₁), we get $\limsup_{n \rightarrow \infty} \tau_n \leq 0$. Now applying Lemma 2.8 to (35), we conclude that $x_n \rightarrow x^*$. Consequently, from $\|J_{r_k, n}^{F_k} x_n - x^*\| \leq \|x_n - x^*\|$, we have $J_{r_k, n}^{F_k} x_n \rightarrow x^*$, for all $k \in \{1, 2, \dots, M\}$.

Corollary 3.2. (see Yao et al. [8]) *Let C be a nonempty closed convex subset of a real Hilbert space H , F a bi-functions from $C \times C$ into \mathbb{R} which satisfy (A₁) - (A₄) and $\{T_i\}_{i=1}^\infty$ an infinite family of nonexpansive mapping of C into C such that $\cap_{i=1}^\infty \text{Fix}(T_i) \cap \text{EP}(F) \neq \emptyset$. Let $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ are three sequences in $(0, 1)$ such that $\alpha_n + \beta_n + \gamma_n = 1$ and $\{r_n\} \subset (0, \infty)$. Suppose the following conditions are satisfied:*

- (B₁) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^\infty \alpha_n = \infty$,
- (B₂) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$,
- (B₃) $\liminf_{n \rightarrow \infty} r_n > 0$ and $\lim_{n \rightarrow \infty} (r_{n+1} - r_n) = 0$.

Let f be a contraction of C into itself with coefficient $\alpha \in (0, 1)$ and given $x_1 \in C$ arbitrarily. Then the sequence $\{x_n\}$ generated by

$$x_{n+1} = \alpha_n f(x_n) + \beta_n x_n + \gamma_n W_n J_{r_n}^F x_n, \quad n \geq 1.$$

converge strongly to $x^ \in \cap_{i=1}^\infty \text{Fix}(T_i) \cap \text{EP}(F)$, where $x^* \in P_{\cap_{i=1}^\infty \text{Fix}(T_i) \cap \text{EP}(F)} f(x^*)$.*

Proof. Take $A = B = 0$, $\phi = \{I\}$, $F_1 = F$ and $F_k = 0$ for $k \in \{2, \dots, M\}$ in Theorem 3.1, then we have $T_{\mu_n} = I$ and $\gamma_n = z_n = J_{r_n}^k x_n$. So from Theorem 3.1 the sequence $\{x_n\}$ converges strongly to $x^* \in \cap_{i=1}^\infty \text{Fix}(T_i) \cap \text{EP}(F)$, where $x^* \in P_{\cap_{i=1}^\infty \text{Fix}(T_i) \cap \text{EP}(F)} f(x^*)$.

Corollary 3.3. *Let C be a nonempty closed convex subset of a real Hilbert space H , $\mathcal{J} = \{F_k : k = 1, 2, \dots, M\}$ be a finite family of bi-functions from $C \times C$ into \mathbb{R} which satisfy (A₁)-(A₄), T a nonexpansive mappings on C such that $\text{Fix}(T) \cap \text{EP}(\mathcal{J}) \neq \emptyset$. Let $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ are three sequences in $(0, 1)$ such that $\alpha_n + \beta_n + \gamma_n = 1$ and $\{r_{k,n}\}_{k=1}^M$ be sequences in $(0, \infty)$. Suppose the following conditions are satisfied:*

- (B₁) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^\infty \alpha_n = \infty$,
- (B₂) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$,
- (B₃) $\liminf_{n \rightarrow \infty} r_{k,n} > 0$ and $\lim_{n \rightarrow \infty} (r_{k,n+1} - r_{k,n}) = 0$ for $k \in \{1, 2, \dots, M\}$.

Let f be a contraction of H into itself and given $x_1 \in H$ arbitrarily. If the sequences $\{x_n\}$ generated iteratively by

$$x_{n+1} = \alpha_n f(x_n) + \beta_n x_n + \gamma_n \frac{1}{n} \sum_{k=0}^\infty \left(\frac{n-1}{n}\right)^k T^k J_{r_{M,n}}^{F_M} \dots J_{r_{2,n}}^{F_2} J_{r_{1,n}}^{F_1} x_n, \quad n \geq 1.$$

Then, sequences $\{x_n\}$ and $\{J_{T_k,n}^{F_k} x_n\}_{k=1}^M$ converge strongly to $x^* \in \text{Fix}(T) \cap \text{EP}(\mathcal{J})$, where $x^* = P_{\text{Fix}(T) \cap \text{EP}(\mathcal{J})} f(x^*)$.

Proof. Let $S = \{0, 1, \dots\}$, $\phi = \{T^i : i \in S\}$ and $T^0 = I$. For $f = (x_0, x_1, \dots) \in B(S)$, define

$$\mu_n(f) = \frac{1}{n} \sum_{k=0}^{\infty} \left(\frac{n-1}{n} \right)^k x_k, \quad \forall n \in \mathbb{N}.$$

Then $\{\mu_n\}$ is a regular sequence of means on $B(S)$ such that $\lim_{n \rightarrow \infty} \|\mu_{n+1} - \mu_n\| = 0$; for more details, see [34]. Next for each $x \in H$ and $n \in \mathbb{N}$, we have

$$T_{\mu_n} x = \frac{1}{n} \sum_{k=0}^{\infty} \left(\frac{n-1}{n} \right)^k T^k x.$$

Take $A = B = 0$, $T_i = I$ for all $i \in \mathbb{N}$ in Theorem 3.1 then we have $y_n = z_n$ and $W_n = I$ for all $n \in \mathbb{N}$. Therefore, it follows from Theorem 3.1 that the sequences $\{x_n\}$ and $\{J_{T_k,n}^{F_k} x_n\}_{k=1}^M$ converge strongly, as $n \rightarrow \infty$ to a point $x^* \in \text{Fix}(T) \cap \text{EP}(\mathcal{J})$, where $x^* = P_{\text{Fix}(T) \cap \text{EP}(\mathcal{J})} f(x^*)$.

Remark 3.4. Theorem 3.1 improve [8, Theorem 1.2] in the following aspects.

(a) Our iterative process (11) is more general than Yao et al. process (14) because it can be applied to solving the problem of finding a common element of the set of solutions of systems of equilibrium problems and systems of variational inequalities.

(b) Our iterative process (11) is very different from Yao et al. process (14) because there are left amenable semigroup of nonexpansive mappings.

(c) Our method of proof is very different from the one in Yao et al. [8] for example we use Corollary 1.1 and Theorem 1.2 of Bruck [33] for the proof of Theorem 3.1.

Acknowledgements

The authors are extremely grateful to the referees for useful suggestions that improved the contents of the paper.

Competing interests

The authors declare that they have no competing interests.

Received: 31 December 2011 Accepted: 16 June 2012 Published: 16 June 2012

References

- Rockafellar, RT: On the maximality of sums of nonlinear monotone operators. *Trans Am Math Soc.* **149**, 75–88 (1970). doi:10.1090/S0002-9947-1970-0282272-5
- Blum, E, Oettli, W: From optimization and variational inequalities to equilibrium problems. *Math Student.* **63**, 123–145 (1994)
- Combettes, PL, Hirstoaga, SA: Equilibrium programming in Hilbert spaces. *J Nonlinear Convex Anal.* **6**, 117–136 (2005)
- Flâm, SD, Antipin, AS: Equilibrium programming using proximal-like algorithms. *Math Program.* **78**, 29–41 (1997)
- Takahashi, S, Takahashi, W: Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces. *J Math Anal Appl.* **331**, 506–515 (2007). doi:10.1016/j.jmaa.2006.08.036
- Bauschke, HH, Borwein, JM: On projection algorithms for solving convex feasibility problems. *SIAM Rev.* **38**, 367–426 (1996). doi:10.1137/S0036144593251710
- Combettes, PL: Quasi-fejérian analysis of some optimization algorithms. *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications* (Haifa, 2000). In *Studies in Computational Mathematics*, vol. 8, pp. 115–152. North-Holland, Amsterdam (2001)
- Yao, Y, Liou, YC, Yao, JC: Convergence theorem for equilibrium problems and fixed point problems of infinite family of nonexpansive mappings. *Fixed Point Theory Appl.* **12** (2007). Article ID 64363
- Jitpeera, T, Katchang, P, Kumam, P: A Viscosity of Ceaàro Mean Approximation Method for a Mixed Equilibrium, Variational Inequality, and Fixed Point Problems **2011**, 24 (2011). Article ID 945051, 838.85. MR 2680251
- Yao, Y, Chen, R, Liou, YC: A unified implicit algorithm for solving the triple-hierarchical constrained optimization problem. *Math Comput Model.* **55**, 1506–1515 (2012). doi:10.1016/j.mcm.2011.10.041
- Yao, Y, Chen, R, Xu, HK: Schemes for finding minimum-norm solutions of variational inequalities. *Nonlinear Anal.* **72**, 3447–3456 (2010). doi:10.1016/j.na.2009.12.029

12. Yao, Y, Cho, YJ, Chen, R: An iterative algorithm for solving fixed point problems, variational inequality problems and mixed equilibrium problems. *Nonlinear Anal.* **71**, 3363–3373 (2009). doi:10.1016/j.na.2009.01.236
13. Yao, Y, Cho, YJ, Liou, YC: Hierarchical convergence of an implicit double-net algorithm for nonexpansive semigroups and variational inequalities. *Fixed Point Theory Appl.* **2011** (2011)
14. Yao, Y, Liou, YC, Kang, SM: Two-step projection methods for a system of variational inequality problems in Banach spaces. *J Glob Optim*
15. Yao, Y, Liou, YC, Marino, G: Strong convergence of two iterative algorithms for nonexpansive mappings in hilbert spaces. *Fixed Point Theory Appl.* (2009)
16. Lau, AT, Miyake, H, Takahashi, W: Approximation of fixed points for amenable semigroups of nonexpansive mappings in Banach spaces. *Nonlinear Anal.* **67**, 1211–1225 (2007). doi:10.1016/j.na.2006.07.008
17. Kangtanyakarn, A: Iterative methods for finding common solution of generalized equilibrium problems and variational inequality problems and fixed point problems of a finite family of nonexpansive mappings. *Fixed Point Theory Appl* **2010**, 29 (2010). Article ID 836714
18. Kim, KS: Convergence of a hybrid algorithm for a reversible semigroup of nonlinear operators in banach spaces. *Nonlinear Anal.* **73**(3), 3413–3419 (2010)
19. Atsushiba, S, Takahashi, W: Approximating common fixed points of nonexpansive semigroups by the Mann iteration process. *Ann Univ Mariae Curie Sklodowska.* **51**, 1–16 (1997)
20. Saeidi, S: Iterative algorithms for finding common solutions of variational inequalities and systems of equilibrium problems and fixed points of families and semigroups of nonexpansive mappings. *Nonlinear Anal.* **70**, 4195–4208 (2009). doi:10.1016/j.na.2008.09.009
21. Piri, H: A general iterative method for finding common solutions of system of equilibrium problems, system of variational inequalities and fixed point problems. *Math Comput Model.* **55**, 1622–1638 (2012). doi:10.1016/j.mcm.2011.10.069
22. Piri, H: Hybrid pseudoviscosity approximation schemes for systems of equilibrium problems and fixed point problems of infinitely and semigroup of nonexpansive mappings. *Nonlinear Anal.* **74**, 6788–6804 (2011). doi:10.1016/j.na.2011.06.056
23. Piri, H: Strong convergence for a minimization problem on solutions of systems of equilibrium problems and common fixed points of an infinite family and semigroup of nonexpansive mappings. *Comput Math Appl.* **61**, 2562–2577 (2011). doi:10.1016/j.camwa.2011.02.049
24. Piri, H, Badali, AH: Strong convergence theorem for amenable semigroups of nonexpansive mappings and variational inequalities. *Fixed Point Theory Appl.* **2011** (2011)
25. Lau, AT, Shioji, N, Takahashi, W: Existence of nonexpansive retractions for amenable semigroups of nonexpansive mappings and nonlinear ergodic theorems in Banach spaces. *J Funct Anal.* **161**(1), 62–75 (1999). doi:10.1006/jfan.1998.3352
26. Agarwal, RP, Donal, O'Regan, Sahu, DR: Fixed point theory for Lipschitzian-type mappings with applications. In *Topological Fixed Point Theory and Its Applications*, vol. 6, Springer, New York (2009)
27. Jitpeera, T, Kumam, P: An extragradient type method for a system of equilibrium problems, variational inequality problems and fixed point of finitely many nonexpansive mappings. *J Nonlinear Anal Optim.* **1**, 71–91 (2010)
28. Suzuki, T: Strong convergence of Krasnoselskii and Mann's type sequences for oneparameter nonexpansive semigroups without Bochner integrals. *J Math Anal Appl.* **305**, 227–239 (2005). doi:10.1016/j.jmaa.2004.11.017
29. Shimoji, K, Takahashi, W: Strong convergence to common fixed points of infinite nonexpansive mappings and applications. *Taiwan J Math.* **5**, 387–404 (2001)
30. Zhang, SS, Joseph Lee, HJ, Chan, CK: Algorithms of common solutions to quasi variational inclusion and fixed point problems. *Appl Math Mech.* **29**, 571–581 (2008). doi:10.1007/s10483-008-0502-y
31. Xu, HK: Viscosity approximation methods for nonexpansive mappings. *J Math Anal Appl.* **298**, 279–291 (2004). doi:10.1016/j.jmaa.2004.04.059
32. Osilike, MO, Igbokwe, DI: Weak and strong convergence theorems for fixed points of pseudocontractions and solutions of monotone type operator equations. *Comput Math Appl.* **40**, 559–567 (2000). doi:10.1016/S0898-1221(00)00179-6
33. Bruck, RE: On the convex approximation property and the asymptotic behavior of nonlinear contractions in Banach spaces. *Israel J Math.* **38**, 304–314 (1981). doi:10.1007/BF02762776
34. Takahashi, W: *Nonlinear Functional Analysis*. Yokohama Publishers, Yokohama (2000)

doi:10.1186/1687-1812-2012-99

Cite this article as: Piri: Approximating fixed points of amenable semigroup and infinite family of nonexpansive mappings and solving systems of variational inequalities and systems of equilibrium problems. *Fixed Point Theory and Applications* 2012 **2012**:99.